وادي الصفوة الزراعية
WADI ALSAFOH

FUNGICIDES

CEERA

Active ingredients:

Azoxystrobin 20%

Difenoconazole 12.5%

Product properties:

A systemic preventive and curative fungicide for a wide range of diseases, including blights, downy mildew, powdery mildew, rots and leaf spots.



Application rates and method

Crop	Target diseases	Application rate	PHI
Cabbage, cauliflower	Anthracnose; leaf spots; powdery mildew	500-750 mL/ha	21 days
Garlic, onion	Purple blotch; downy mildew; Alternaria blight	500-750 mL/ha	21 days
Carrot	Leaf blight; powdery mildew	500-750 mL/ha	14 days
Cucumber, zucchini, watermelon, melon	Leaf blight; anthracnose; downy mildew; powdery mildew	50-70 mL/100 L water	3 days
Grape	Early blight; powdery mildew; black rot; downy mildew; fruit rot	50-70 mL/100 L water	21 days
Wheat	Powdery mildew; rust	500-750 mL/ha	30 days
Pepper	Powdery mildew; leaf spot	50-70 mL/100 L water	3 days
Potato	Early blight	500-750 mL/ha	14 days
Tomato	Early blight; powdery mildew	50-70 mL/100 L water	7 days