وادي الصفوة الزراعية
WADI ALSAFOH

FUNGICIDES

EMOTION

Active ingredient:

Propamocarb hydrochloride 72.2%

Product properties:

A systemic preventive and curative fungicide effective against downy mildew and root rot. Rapid absorption and translocation provide uniform protection of new growth. It prevents fungal spore germination, stops mycelial development and provides long-lasting protection.



Application rates and method

Crop	Target diseases	Application rate	PHI
Vegetables: sweet pepper, tomato, onion, lettuce, spinach, cucumber	Root rot: Pythium, Phytophthora; downy mildew	Seed treatment: after sowing, spray the soil with 100 mL/100 L water at 5 L solution/m2. Seedlings: 2-4 days before planting, use the same soil rate, then transplant. Five days after transplanting, use 100 mL/100 L water at 100-150 mL solution/plant; repeat every 2 weeks if needed. Foliar spray: 150-300 mL/100 L water; repeat every 7-10 days at early infection. Maximum 3 sprays/season.	14 days; lettuce: 21 days; cucumber: 4 days
Ornamental plants	Root rot: Pythium, Phytophthora; downy mildew	Seeds and pre-transplant soil: 5-10 mL/m2 in 1-5 L water depending on soil moisture. Cuttings before planting: 10-15 mL/m2 in 1-5 L water.	-